

COMPLETE HYPERBOLIC STEIN MANIFOLDS WITH PRESCRIBED AUTOMORPHISM GROUPS

SU-JEN KAN

ABSTRACT.

It is well-known that the automorphism group of a hyperbolic manifold is a Lie group. Conversely, it is interesting to see whether or not any Lie group could be prescribed as the automorphism group of certain complex manifold.

When the Lie group G is compact and connected, this problem has been completely solved by Bedford-Dadok and independently by Saerens-Zame on 1987. They have constructed strictly pseudoconvex bounded domains Ω such that $Aut(\Omega) = G$. For Bedford-Dadok's Ω , $0 \leq \dim_{\mathbb{C}}\Omega - \dim_{\mathbb{R}}G \leq 1$; for generic Saerens-Zame's Ω , $\dim_{\mathbb{C}}\Omega \gg \dim_{\mathbb{R}}G$.

J. Winkelmann has answered affirmatively to noncompact connected Lie groups in recent years. He showed there exist Stein complete hyperbolic manifolds Ω such that $Aut(\Omega) = G$. In his construction, it is typical that $\dim_{\mathbb{C}}\Omega \gg \dim_{\mathbb{R}}G$.

In this article, we tackle this problem from a different aspect. We prove that for any connected Lie group G (compact or noncompact), there exist complete hyperbolic Stein manifolds Ω such that $Aut(\Omega) = G$ with $\dim_{\mathbb{C}}\Omega = \dim_{\mathbb{R}}G$. Working on a natural complexification of the real-analytic manifold G , our construction of Ω is geometrically concrete and elementary in nature.

1. Introduction.

It is well-known that the automorphism group of a hyperbolic manifold has the structure of a Lie group. It is natural to ask whether every Lie group could appear as the automorphism group of certain hyperbolic manifolds or not. This problem sometimes is called the realization problem: realizing a Lie group as the automorphism group of certain complex manifold.

This realization problem has been completely solved by Bedford-Dadok [B-D] and independently by Saerens-Zame [S-Z] when the given Lie group is compact and connected. In recent years, generalizing ideas of Saerens-Zame, Winkelmann [W] has settled the realizing problem for any connected Lie group. However, domains Ω constructed from this Saerens-Zame-Winkelmann's approach are typically with $\dim_{\mathbb{C}}\Omega \gg \dim_{\mathbb{R}}G$.

2000 *Mathematics Subject Classification.* 32C09, 32Q28, 32Q45, 53C24, 58D19. This research is partially supported by NSC 93-2115-M-001-006.

The strategy of Saerens-Zame-Winkelmann was first to find a domain D on which G acts by automorphisms, and then perturbed it to a G -invariant strictly pseudoconvex subdomain in such a way that the additional automorphisms were ruled out by assigning CR-invariants to each G -orbit on the boundary. To find such a domain D to start with, Saerens-Zame first embedded the compact Lie group G into the unitary group $U(N_1)$ and then constructed a domain D in $GL(N_1, \mathbb{C}) \times \mathbb{C}^{N_2}$ on which G acts by automorphisms where N_1 and N_2 are large in general. Due to the above embedding process, the generic resultant complex manifold Ω has large complex dimension, $\dim_{\mathbb{C}} \Omega \gg \dim_{\mathbb{R}} G$. Having observed every Lie algebra is linear and hence the universal covering of a Lie group could be viewed as linear, Winkelmann has been able to embed $\tilde{G} \hookrightarrow Sp(N_3, \mathbb{R})$ and then find a suitable domain $D \subset \mathbb{C}^{N_4}$ to start with. Again, this embedding process and the construction of the domain D has enormously increased the dimension.

The complexification $G_{\mathbb{C}}$ of a compact Lie group G is Stein with $\dim_{\mathbb{C}} G_{\mathbb{C}} = \dim_{\mathbb{R}} G$. Starting from domains in $G_{\mathbb{C}}$, Bedford-Dadok were able to give a more concrete construction. They found bounded strictly pseudoconvex domains $\Omega \subset G_{\mathbb{C}}$ or $\Omega \subset G_{\mathbb{C}} \times \mathbb{C}$ such that $Aut(\Omega) = G$.

A natural attempt is to generalize Bedford-Dadok's approach to noncompact Lie groups. Unfortunately, it is not easy to give a canonical complexification of a generic noncompact Lie group such that $\dim_{\mathbb{C}} G_{\mathbb{C}} = \dim_{\mathbb{R}} G$. In this article we consider a special kind of complexifications of real-analytic manifolds to resolve this difficulty.

A Lie group G equipped with a left-invariant metric g is naturally a homogeneous space with $Isom(G) \simeq L(G) \cdot K$ where K is the isotropy group and $L(G)$ are left multiplications. We provide the tangent bundle of the Riemannian manifold (G, g) with a complex structure in such a way that all the leaves of the Riemannian foliation are holomorphic curves. The disk bundle of radius r is called a Grauert tube $T^r G$.

Generalizing the rigidity result, $Aut(T^r G) = Isom(G)$, proved in [K] we are able to dominate automorphism groups of certain domains $D \subset T^r G$. We prove it at Theorem 3.5 and call it the subrigidity $Aut(D) < Isom(G)$.

Starting from such a domain D our strategy is to destroy, through certain G -invariant perturbations, the additional automorphisms coming from the isotropy group. Though the objects we deal with are not even relatively compact, the holomorphic extension property needed here is not hard to handle due to the special

complex structure adopted here. In fact, most mappings we deal with are bundle mappings which automatically extend over the boundary. The main result of this article is

Theorem. *Let G be a connected Lie group of dimension $n \geq 2$. Then there exist complete hyperbolic Stein manifolds Ω , $\dim_{\mathbb{C}}\Omega = n$, such that $\text{Aut}(\Omega) = G$.*

The dimensional condition $n \geq 2$ has to be added here since the main idea we'll use follows from the rigidity arguments of Grauert tubes while Grauert tubes and their perturbations are biholomorphic to the unit disc, by the Riemann mapping theorem, when the center is of dimension one. In the rigidity arguments of Grauert tubes, it is also necessary to assume the Riemannian manifold is connected. At this moment, I could not see a way to release the connectedness of the Lie groups. We remark while Saerens-Zame's result works for compact disconnected Lie group as well, the resultant complex manifold Ω is typically with $\dim_{\mathbb{C}}\Omega \gg \dim_{\mathbb{R}}G$.

We emphasize that methods developed in this article work simultaneously for compact and noncompact Lie groups; the complex dimension of the resultant complex manifold Ω is equal to the real dimension of G ; and Ω thus obtained are constructed in a geometrically concrete and nature way.

The organization of this article is the following. In §2 we briefly review terminologies concerning Grauert tubes and prove the existence of Stein Grauert tubes. Generalizing notations about the rigidity of Grauert tubes, we prove subrigidity characterization of certain domains in §3. In §4, specific domains and perturbations are constructed explicitly. We perturb domains in the tangent bundle of a connected Lie group in an invariant way such that extra symmetry on each fiber would be eliminated. By constructed such kind of domains in a fairly explicitly way, the realization of a connected Lie group as an automorphism group follows from the subrigidity derived on §3.

This realization problem was first noticed by the author and also mentioned to the author by Peter Heinzner in September 2002 when I was visiting Ruhr-Universität, Bochum Germany. Over years I was trapped in an effort to show every Grauert tube is Stein. Among other things, I am very grateful to Peter Heinzner for bringing my attention to [HHK] and [D-G], which have helped settling down the Steiness part needed in this article, during my recent visit to Bochum in August 2004. I would also like to thank Jörg Winkelmann for many helpful discussions about this project during the Hayama Symposium 2002.

2. Existence of Stein Grauert tubes.

Throughout this article, (M, g) is assumed to be a connected real-analytic Riemannian manifold of dimension ≥ 2 . There exists a complex structure, the *adapted complex structure*, in a domain $\Omega(M) \subset TM$ which turns leaves of the Riemannian foliation on $\Omega(M) \setminus M$ into holomorphic curves. Let's denote by $\Omega(M)$ the maximal domain in which the adapted complex structure exists. With respect to this complex structure, the length square function $\rho(x, v) := |v|^2$ is strictly plurisubharmonic and satisfies the complex homogeneous Monge-Ampère equation $(dd^c \rho)^n = 0$ on $\Omega(M) \setminus M$ with the initial condition $\rho_{i\bar{j}}|_M = \frac{1}{2}g_{ij}$. The *Grauert tube*

$$T^r M = \{(x, v) : x \in M, v \in T_x M, |v| < r\} = \{\rho^{-1}[0, r^2]\}$$

is the collection of tangent vectors of length less than r equipped with the adapted complex structure; M is the center and r is the radius. In general the maximal domain $\Omega(M)$ is not a Grauert tube, $\Omega(M)$ is a Grauert tube if and only if M is a symmetric space of rank one. There is a natural antiholomorphic involution σ fixing every point of M ,

$$(2.1) \quad \sigma : \Omega(M) \rightarrow \Omega(M), \quad (x, v) \rightarrow (x, -v).$$

For two isometric real-analytic Riemannian manifolds (M, g) and (N, κ) , the nature of the adapted complex structure will assure the biholomorphic equivalence of $\Omega(M)$ and $\Omega(N)$ and the biholomorphic equivalence of $T^r M$ and $T^r N$ as well. Given an isometry h of (M, g) , the differential dh acts as a biholomorphism on $\Omega(M)$;

$$(2.2) \quad dh : \Omega(M) \rightarrow \Omega(M), \quad dh(x, v) = (h(x), h_* v).$$

The notation F_p^r is reserved for the fiber passing $p \in M$,

$$(2.3) \quad F_p^r := \{(p, v) : v \in T_p M, |v| < r\}.$$

Since the complex structure we consider here is a local object, for each real-analytic Riemannian manifold, there exists an $r_{max}(M) \geq 0$, the maximal radius such that the adapted complex structure could be well-defined on $T^{r_{max}} M$. For compact, co-compact or homogeneous M , $r_{max}(M) > 0$. For any $r < r_{max}(M)$, the

complex manifold $T^r M \subset T^{r_{max}} M$ with strictly pseudoconvex boundary defined by $\{\rho = r^2\}$.

It is shown in Theorem 5.2 of [K] that any Grauert tube over homogeneous space of radius $r < r_{max}$ is complete hyperbolic. Another natural question to ask about is whether or not a Grauert tube is Stein.

Recall from [D-G] that an *unramified Riemannian domain* D over a Stein manifold X is a complex manifold D together with a locally homeomorphic holomorphic map from D to X and that a pseudoconvex unramified Riemannian domain over a Stein manifold is itself Stein. It is clear that any pseudoconvex domain in a Stein manifold is Stein since we may take the imbedding as the holomorphic map.

Proposition 2.1. *$T^r M$ is a Stein manifold for all $r \leq r_{max}(M)$ if M is one of the following: compact; co-compact; homogeneous with nonnegative curvature.*

Proof. For compact M , the Grauert tube $T^r M, r \leq r_{max}$ is exhausted by the strictly plurisubharmonic function $-\log(r^2 - \rho)$, hence is Stein.

When M is co-compact, $\hat{M} = M/\Gamma$ is compact for some discrete subgroup $\Gamma < Isom(M)$, then the Grauert tube $T^r M$ is simply a covering of the Stein Grauert tube $T^r \hat{M}$. Thus $T^r M, r \leq r_{max}(M)$, is Stein.

Following the decomposition theorem (Theorem 7.1*, [C-G]) of Cheeger-Gromoll a homogeneous manifold M of nonnegative curvature may be written as the product $\mathbb{R}^k \times M^*$ where M^* is a compact homogeneous space of nonnegative curvature. Since the metric on M is the product metric from $\mathbb{R}^k$ and M^* , it is clear that $r_{max}(M) = r_{max}(M^*)$ and $T^r(M) \subset \mathbb{C}^k \times T^{r_{max}}(M^*)$, $\forall r \leq r_{max}$. For any $r \leq r_{max}$, $T^r(M)$ has pseudoconvex boundary since the boundary is locally exhausted by the strictly plurisubharmonic function $-\log(r^2 - \rho)$. Being a pseudoconvex domain in the Stein manifold $\mathbb{C}^k \times T^{r_{max}}(M^*)$, $T^{r_{max}} M$ is Stein. $\square$

Though there is no example disproving the Steiness of any Grauert tube, the above three categories are the only complete classes that we are sure about the Steiness of any Grauert tubes over. For some very special kind of homogeneous spaces $M = G/K$, Halverscheid-Iannuzzi [H-I] were able to construct a polar map from TM to the Stein manifold $G_{\mathbb{C}}/i(K_{\mathbb{C}})$. They show the maximal domain over a Heisenberg group is neither holomorphically separable nor holomorphically convex. However, there are some facts hiding behind: Grauert tubes constructed over their spaces are Stein. Their polar map is an unramified one therefore, Grauert tubes are pseudoconvex Riemann domains over the Stein manifold $G_{\mathbb{C}}/i(K_{\mathbb{C}})$, they are

Stein (c.f. [D-G]).

A complex manifold X is a *complexification* of M if $M \subset X$ as a maximal totally real submanifold $\dim_{\mathbb{C}} X = \dim_{\mathbb{R}} M$. Utilizing his solution to the Levi problem, Grauert has proved there always exists a Stein complexification of a real-analytic manifold.

Though any two complexifications (X_1, J_1) and (X_2, J_2) of a real-analytic manifold M are locally biholomorphic near M , i.e., there exist a neighborhood U_1 of M in X_1 and a neighborhood U_2 of M in X_2 such that U_1 and U_2 are biholomorphically equivalent, we still can't conclude there are some Stein Grauert tubes of small radii since it does not seem clear how to control the radius uniformly when the manifold is not relatively compact.

In [HHK], the authors have put on an extra piece of information, say there is some kind of G -action on M . They ask for the question whether the Stein manifold in Grauert's complexification could still be chosen to be G -invariant or not. Analyzing the real-analytic slices of certain categorical quotients, they conclude the following:

Theorem(H-H-K). *Let X be a G -complexification of a real-analytic manifold M where G is a connected Lie group acting properly on M as real-analytic diffeomorphisms. Then there exists a G -invariant Stein neighborhood of M in X .*

A group action G on M is *proper* if the inverse image of a compact set is compact. A complexification X of M is called a *G -complexification* if the G -action on M extends to a holomorphic G -action on X .

Though it is not clear whether a generic Grauert tube is Stein or not, a direct application of the above theorem would assert the existence of Stein Grauert tubes of small radii provided the centers have possessed some transitivity property. This would be enough for the purpose of solving the realization problem.

Proposition 2.2. *Let (M, g) be a homogeneous space. Then there exist $\epsilon > 0$ such that the Grauert tube $T^\epsilon M$ is Stein and complete hyperbolic.*

Proof. Let $G = \text{Isom}_0(M)$, then G acts properly on M and acts holomorphically on the complexification $T^{r_{max}} M$ of M . By [HHK], there exists a G -invariant Stein neighborhood U of M in $T^{r_{max}} M$. For $p \in M$ we may assume the fiber F_p^ϵ , tangent vectors at p of length less than ϵ , is contained in U . By the transitivity of the G -action, $G \cdot F_p^\epsilon$ is the Grauert tube $T^\epsilon M$. The Steinness follows since $T^\epsilon M$ is now a strictly pseudoconvex domain in the Stein manifold U . $\square$

3. Subrigidity of domains.

Let Aut denote the automorphism group and $Isom$ denote the isometry group; Aut_0 and $Isom_0$ denote the corresponding identity components. For any real-analytic Riemannian manifold (M, g) such that $T^r M$ is not covered by the ball, the following rigidity results were proved in [K].

- (1) $Aut_0(T^r M) = Isom_0(M)$;
- (2) furthermore, if M is homogeneous then $Aut(T^r M) = Isom(M)$.

We remark here that whenever the rigidity is mentioned, we always assume $\dim M > 1$. When $\dim M = 1$, the Grauert tube $T^r M$ is the disc in $\mathbb{C}$. One essential feature of Grauert tubes is the symmetry on each fiber, each fiber is a disk bundle. In this section, we'll consider domains in a more general setting, perfect symmetry on fibers won't be asked for any more. One crucial step in proving the rigidity is the Theorem 4.1 in [K] which has characterized the isometry group of M . We will show this kind of characterization still works and $Aut(D)$ is dominated by $Isom(M)$ provided there is some transitivity on M . Arguments will follow the spirits and methods developed in [K]. However, an extra piece of assumptions on the target Grauert tube is necessary; we assume Grauert tubes are Stein while the existence has been guaranteed in the last section.

Lemma 3.1. *Let $T^r M$ be a Stein Grauert tube, $D \subset T^r M$ be a connected domain containing M . If $f \in Aut(D)$, $f(M) = M$, then $f = du$ for some isometry u from (M, g) to (M, κ) and f could be extended over the boundary ∂D .*

Proof. From the construction of Grauert tubes, it is clear that M is a maximal totally real submanifold of D . Let ρ denote the length square function on $T^r M$, then $\rho \cdot f^{-1}$ is a strictly plurisubharmonic function on D . The Riemannian metric g is induced from the Kähler form $i\partial\bar{\partial}\rho$, $2\rho_{i\bar{j}}|_M = g_{ij}$. Similarly, the Kähler form $i\partial\bar{\partial}(\rho \cdot f^{-1})$ has induced a Riemannian metric k on M . Let $u = f|_M : (M, g) \rightarrow (M, \kappa)$; u is an isometry. By the nature of the adapted complex structure, the differential du is a biholomorphic mapping from the Grauert tube $T^r(M, g)$ to the Grauert tube $T^r(M, \kappa)$.

Let $D_1 = D \cap T^r(M, \kappa) \hookrightarrow \mathbb{C}^N$ for some N . This could be achieved since $T^r(M, g)$ is Stein and $T^r(M, \kappa)$ is biholomorphic to $T^r(M, g)$. Let $D_2 = (du)^{-1}(D_1) \subset T^r(M, g)$. Let $D_3 = D \cap D_2$ then D_3 is a domain in $T^r M$ which has contained M as a maximal totally real submanifold. Restricted to D_3 , f and du could be viewed

as holomorphic functions from D_3 to $\mathbb{C}^N$. Since $f = du$ at the maximal totally real submanifold M , by the identity principle, $f \equiv du$ on the subdomain D_3 and hence $f \equiv du$ on D . For $(x, v) \in D$, $f(x, v) = du(x, v) = (u(x), u_*v) = (f(x), u_*v)$ is a bundle map acting fiberwise. It is clear that f could be extended over the boundary of D . $\square$

Recall $\Omega(M)$ is the maximal domain in TM such that the adapted complex structure is defined; σ is the natural antiholomorphic involution in $\Omega(M)$ and the norm-square function ρ is strictly plurisubharmonic in $\Omega(M)$. Sibony (Theorem 3, [S]) has asserted the hyperbolicity of any complex manifold equipped with a bounded strictly plurisubharmonic function. For domains in $\Omega(M)$ to be hyperbolic, the only thing we got to take care is the vertical direction.

One essential feature of Grauert tubes is that each fiber, which is diffeomorphic to a real ball, has perfect symmetry. We have tried to release this symmetry on the fiber to get some kind of rigidity result. However, it seems to us certain kind of transitivity on the domain is necessary for our purpose. We make the following definition.

Definition 3.2. *A domain $D \subset \Omega(M)$ is G -homogeneous if:*

- (1) *there exists a connected subgroup $G \subset \text{Isom}(M)$ acting transitively on M ;*
- (2) *there exists a bounded open subset $F_p \subset T_p M$ such that $D = G \cdot F_p$.*

With this extra piece of symmetry, Lemma 3.1 has the following refined form:

Proposition 3.3.

Let D be a G -homogeneous domain in $\Omega(M)$. If $f \in \text{Aut}(D)$, $f(M) = M$, then $f = du$ for some $u \in \text{Isom}(M)$.

Proof. Since M is homogeneous, there exists small ϵ such that $T^\epsilon M$ is Stein. The restriction map of f to $D \cap T^\epsilon M$ has possessed the bundle map property developed in Lemma 3.1. Thus $f = du$, for some isometry u from the Riemannian manifold (M, g) to the Riemannian manifold (M, κ) . By the assumption on D there exist $G \subset \text{Isom}(M)$ and $p \in M$ such that $D = G \cdot F_p$.

Let $q = u(p) = h(p)$ for some $h \in G$ and let $\{e_1, \dots, e_n\}$ be an orthonormal basis of F_p with respect to the metric g . Since h is an isometry, $\{(e_1, \dots, e_n)A\}$ is an orthonormal basis of F_q with respect to the metric g where $A \in O(n)$. On the other hand, $\{u_*e_1, \dots, u_*e_n\}$ form an orthonormal basis of F_q with respect to the

metric κ , thus, there exists a $B \in GL(n, \mathbb{R})$ such that

$$(u_*e_1, \dots, u_*e_n) = (e_1, \dots, e_n)AB.$$

Let's denote the matrix $AB = C = (C_1, \dots, C_n)$ and $e = (e_1, \dots, e_n)$ then

$$\delta_{ij} = \langle u_*e_i, u_*e_j \rangle_\kappa = \langle eC_i, eC_j \rangle_g = C_i \cdot C_j.$$

This shows C is a matrix with orthonormal columns, thus $C \in O(n)$ and $B \in O(n)$ as well.

The orthonormal basis $\{u_*e_1, \dots, u_*e_n\}$ of the metric κ come from an orthogonal transformation of the orthonormal basis $\{e_1, \dots, e_n\}$ of the metric g . Thus, $\kappa = g$ and $u \in Isom(M, g)$. $\square$

A G -homogeneous domain is hyperbolic and hence its automorphism group is a Lie group. Let D be a σ -invariant strictly pseudoconvex G -homogeneous domain in $\Omega(M)$. Following notations used in §6 of [K], we set $\widehat{Aut}(D) := Aut(D) \cup \sigma \cdot Aut(D)$ and denote $\hat{\mathcal{L}}$ as the Lie algebra of $\widehat{Aut}(D)$. Modifying the proofs of Lemma 6.1, Lemma 6.2 and Lemma 6.3 in [K], it could be shown that for every $\xi \in \mathcal{L}$, the Lie algebra of $Aut(D)$, the vector field associated to ξ is tangent to M . We conclude

Proposition 3.4. *Let D be a σ -invariant strictly pseudoconvex G -homogeneous domain in $\Omega(M)$. If D is not biholomorphic to the ball. Then $Aut_0(D) \subset Isom_0(M)$.*

Furthermore, following [K] again, we are going to prove the subrigidity $Aut(D) \subset Isom(M)$. For the rest of this section, we assume D is a strictly pseudoconvex σ -invariant G -homogeneous domain in $\Omega(M)$, D is not biholomorphic to the ball. Proofs go exactly the same way as in §7 of [K]; a brief explanation would be given in the following.

(I). First of all, a strictly pseudoconvex G -homogeneous domain in $\Omega(M)$ is complete hyperbolic since the boundary behavior is dominated by the strictly pseudoconvexity and the horizontal direction is determined by the transitivity of the G -action.

(II). Secondly, the following is clear from the G -homogeneity and Prop. 3.4.

$$G \subset Aut_0(D) \subset Isom_0(M).$$

Given $f \in Aut(D)$, D is also a $f \cdot G \cdot f^{-1}$ -homogeneous domain centered at $N = f(M)$. By the homogeneity of the G -action,

$$N = f(M) = f(G \cdot p) = f(Aut_0(D) \cdot p) = Aut_0(D) \cdot f(p);$$

$$\pi(N) = \pi(Aut_0(D) \cdot f(p)) = Aut_0(D) \cdot \pi(f(p)) = M.$$

Thus, the argument in Lemma 7.1 of [K] goes through and $f(M) \cap T_p M \neq \emptyset$ for any $p \in M$.

(III). Since D is complete hyperbolic and G -homogeneous, arguments in Prop.7.2 of [K] could be transplanted here. Thus, the index of G in $Aut(D)$ is finite.

(IV). Let $\tau = f \cdot \sigma \cdot f^{-1}$ be the associated antiholomorphic involution with respect to the fixed point set $N = f(M)$, then the identity relation at (7.2) and the Proposition 7.3 of [K] work as well. Thus, there exists an odd integer k such that $(\sigma \cdot \tau)^k = id$.

We conclude this section by the following theorem.

Theorem 3.5.

Let D be a σ -invariant strictly pseudoconvex G -homogeneous domain in $\Omega(M)$. Then either $Aut(D) \subset Isom(M)$ or D is the ball.

Proof. As stated in (III), $Aut(D)/G = \{g_j G : g_j \in Aut(D), j = 1, \dots, k\}$. Then $\psi(z) = \sum_{j=1}^k \rho(g_j(z))$ is an $\widehat{Aut}(D)$ -invariant strictly plurisubharmonic nonnegative function in D where ρ is the length square function in D . As G acts transitively on M , the tangent space $T_z(D)$ could be decomposed as, for any $z \in D$,

$$T_z(D) = T_z(G \cdot z) + T_z(T_{\pi(z)}M \cap D)$$

where π is the fiber projection $\pi(x, v) = x, \forall v \in T_x M$.

Since ψ is constant in $G \cdot z$, every critical point of the function $f_z := \psi|_{T_{\pi(z)}M \cap D}$ is a critical point of ψ and every critical point of ψ occurs at the critical points of the functions f_z .

As ψ is strictly plurisubharmonic, the above decomposition implies that the real Hessian of f_z is positive definite on the tangent space $T_z(T_{\pi(z)}M \cap D)$. Since f_z is proper on the fiber, it follows that there is exactly one critical point of f_z which turns out to be the minimal point. Since $\psi \cdot \sigma = \psi$, the minimum of f_z occurs at $\pi(z)$. That is to say that the set of critical points of ψ is exactly M .

Let $f \in Aut(D)$, $f(M) = N$, then N is the critical point set of ψ since ψ is $Aut(D)$ -invariant. We conclude that $N = M$ and $f \in Isom(M)$ following from Prop.3.3. The inclusion $Aut(D) \subset Isom(M)$ is concluded. $\square$

4. Realizing a connected Lie group as an automorphism group.

Let G be a connected real Lie group with Lie algebra $\mathcal{G}$. Given a positive definite inner product $\langle \cdot, \cdot \rangle$ on $T_e G = \mathcal{G}$, we may endow G with the associated left

invariant Riemannian metric g . Every Lie group is real-analytic, since it is locally diffeomorphic to the Lie algebra $\mathcal{G}$ through the exponential map. Thus (G, g) is a real-analytic Riemannian manifold. Furthermore, (G, g) is a homogeneous space with trivial tangent bundle $G \times T_e G$.

It is clear that the left translation group $L(G)$ is a subgroup of $Isom(G, g)$ and $L(G)$ acts transitively on G . Furthermore, the transitivity implies the diffeomorphic equivalence of G with $Isom(G, g)/K$ and with $Isom_0(G, g)/K_0$ where

$$(4.1) \quad K = \{h \in Isom(G, g) : h \cdot e = e\}$$

is the isotropy group at $e \in G$ and $K_0 = K \cap Isom_0(G, g)$. The following two equations are immediate.

$$(4.2) \quad Isom(G, g) \simeq L(G) \cdot K, \quad Isom_0(G, g) \simeq L(G) \cdot K_0.$$

From now on, we denote the homogeneous Riemannian manifold (G, g) as (M, g) where G is a connected Lie group of dimension $n \geq 2$ and g is a left invariant real-analytic metric on G . We would like to construct a G -homogeneous strictly pseudoconvex domain and then further perturb the domain to eliminate additional automorphisms such that the automorphism group of the resultant domain is G . We emphasize that our method work for both compact and noncompact Lie groups as long as the Lie group is connected. The dimensional condition $n \geq 2$ has to be added here since the main idea we used here follows from the rigidity argument of Grauert tubes while the rigidity of Grauert tubes fails when the center is of dimension one.

Let $(M, g) = (G, g)$ be as above. Since it is homogeneous, there exist $\delta > 0$ such that the Grauert tube $T^\delta M$ is Stein and the Grauert tube $T^{2\delta} M$ still exists. Let ρ denote the length square function; σ denote the natural antiholomorphic map in $T^{2\delta} M$ and F_p^r denote tangent vectors at $p \in M$ of length less than r , $F_p^r := \{(p, v) : v \in T_p M, |v| < r\}$.

Let $\{e_1, \dots, e_n\}$ be an orthonormal basis—with respect to the metric g —of $F_p^{2\delta}$, K be the isotropy group of (M, g) at $p \in M$. Then $K \subset O(n)$ when we view $F_p^{2\delta}$ as a subspace of the real vector space generated by the orthonormal basis $\{e_1, \dots, e_n\}$.

Let $a_j = \delta e_j \in F_p^{2\delta}, j = 1, \dots, n$. Choose a ball $B_\epsilon \subset F_p^{2\delta}$ centered at a_1 of radius $\epsilon \ll \delta$ and a family of orthogonal transformations $f_j \in O(n)$ such that $f_j(a_1) = a_j, j = 2, \dots, n$.

Let $\eta_1(x) = (x - a_1)^{4l}$, $l \geq 1$, be a real-analytic function on B_ϵ . Convoluting with some cut-off function, we may assume η_1 has compact support $C_1 \subset B_\epsilon$.

Denoting $f_1 = id$, we define a function η_2 on $F_p^{2\delta}$ as follows:

$$(4.3) \quad \eta_2(f_j(x)) = \eta_2(\sigma(f_j(x))) = \frac{1}{j}\eta_1(x), \quad \forall x \in F_p^{2\delta}, j = 1, \dots, n.$$

The ball B_ϵ could be arranged so small that $\{\sigma^i(f_j(B_\epsilon)) : i = 0, 1; j = 1, \dots, n\}$ are pairwise disjoint. It is clear that η_2 is a σ -invariant function in $F_p^{2\delta}$ with compact support C_2 contained in $\cup_{j=1}^n (f_j(B_\epsilon) \cup \sigma(f_j(B_\epsilon)))$.

Since the tangent bundle of a Lie group is trivial, $TM = G \times T_p M$. We define a real-analytic functions η in $T^{2\delta}M = G \times F_p^{2\delta}$ by setting

$$(4.4) \quad \eta(h \cdot x) = \eta_2(x), \quad \forall h \in G, x \in F_p^{2\delta}.$$

Clearly, η is a nonnegative G -invariant and σ -invariant real-analytic function on $T^{2\delta}M$. Recall ρ is the length square function of the tangent vectors which is G -invariant and strictly plurisubharmonic and $T^\delta M = \{\rho^{-1}([0, \delta^2])\}$. Define the function, for some $0 < \epsilon' \ll 1$,

$$(4.5) \quad \hat{\rho} = \rho - \delta^2 + \epsilon' \eta$$

Shrinking ϵ' if necessary such that all the 0-th, first and second order derivatives of $\epsilon' \eta$ are well under control. Thus, we may assume the function $\hat{\rho}$ is strictly plurisubharmonic. It is G and σ invariant since both ρ and η are.

Let D be a domain defined by the function $\hat{\rho}$,

$$(4.6) \quad D := \{z \in T^{2\delta}M; \hat{\rho}(z) < 0\}.$$

It is clear that D is a σ -invariant strictly pseudoconvex G -homogeneous domain in $T^\delta M$. In order to apply Theorem 3.5 to conclude the subrigidity, we need to show the domain D is not biholomorphic to the ball. Some background on Chern-Moser normal form is needed here.

In the fundamental paper [C-M], Chern and Moser have associated to every strictly pseudoconvex point p in a hypersurface H a family of local invariants, namely a neighborhood U_p of p in H is biholomorphically equivalent to a neighborhood V_q of q in a hypersurface S if and only if the associated families of invariants at p and at q are the same. These invariants are given by the coefficients of certain normal form of the defining function which we briefly explain in the following. Let

ψ be a local defining function of the $(2n-1)$ -dimensional hypersurface H near the point p , say inside a coordinate chart. Since p is strictly pseudoconvex the Levi form is positive definite and the defining function could be transformed through some linear translation and proper holomorphic transformations to the following:

$$(4.7) \quad v = |z|^2 + F(z, \bar{z}, w)$$

where $z = (z_1, \dots, z_{n-1}) \in \mathbb{C}^{n-1}$, $w = u + iv \in \mathbb{C}$. Using the transformations $z^* = z + f(z, w)$, $w^* = w + g(z, w)$ Chern and Moser have simplified (4.7) to

$$(4.8) \quad v^* = |z^*|^2 + N_{24}(u^*)z^{*2}\bar{z}^{*4} + N_{42}(u^*)z^{*4}\bar{z}^{*2} + \sum_{j,k \geq 2, j+k \geq 7} N_{jk}(u^*)z^{*j}\bar{z}^{*k}$$

Furthermore this transformation, and hence all of the coefficients N_{jk} , is made unique when certain normalizations on f and g are made.

For the hyperquadric $Q = \{v = |z|^2\}$, the unbounded model of the sphere, all invariants N_{jk} vanish.

Lemma 4.1. *D is not biholomorphic to the ball.*

Proof. It is standard that given any n -dimensional totally real closed submanifold $X \subset B^n$ there exists $\varphi \in \text{Aut}(B^n)$ such that $\varphi(X) = \mathbb{R}^n \cap B^n$.

Suppose there exists a biholomorphic map $f : D \rightarrow B^n$, then $f(M)$ is a maximal totally real closed submanifold of B^n since M is such kind of submanifold of D . Without loss of generality, we may assume $f(M) = \mathbb{R}^n \cap B^n$. Adopting the argument used in the proof of Lemma 3.1, we see f is a bundle map, $f = du$ for some isometry u from (M, g) to $(\mathbb{R}^n \cap B^n, g^*)$.

Thus, the biholomorphic map f could be extended holomorphically over the boundary ∂D and thus the Chern-Moser normal form of $\hat{\rho}$ at any boundary point of D has the local expression $v = |z|^2$.

However, the defining function near $a_1 \in \partial D$ and the defining function near generic boundary points differ by some $4l$ -order terms. In the construction of the normal form, we see the orders won't decrease. Thus the Chern-Moser normal form at a_1 won't be the same as the Chern-Moser normal form at generic boundary points which we assume to be $v = |z|^2$. A contradiction. $\square$

It is clear G is contained in the automorphism group of D since D is G -invariant. Theorem 3.5 along with (4.2) implies

$$(4.9) \quad L(G) < \text{Aut}(D) < \text{Isom}(M) \simeq L(G) \cdot K.$$

The last step is to eliminate those automorphisms coming from the isotropy group.

Theorem 4.2. *D is a complete hyperbolic Stein manifold with $\text{Aut}(D) = G$.*

Proof. D is complete hyperbolic since it is G -homogeneous with strictly pseudoconvex boundary. It is Stein since D is a strictly pseudoconvex domain in the Stein manifold $T^\delta M$.

It remains to show if $h \in K \cap \text{Aut}(D)$, then h is the identity map. By the construction of η_2 at (4.3) and hence the construction of the defining function $\hat{\rho}$, we see that there exist neighborhoods U_j of a_j on the hypersurface $\partial D \cap F_p^\delta$ such that points in $U_i - a_i$ all have norms different from norms of points in $U_j - a_j$ for $i \neq j$. Besides, every point in $U_i - a_i$ has norm $< \delta$ and $|a_i| = \delta$, $i = 1, \dots, n$.

Since K is a subgroup of $\text{Isom}(M)$, h is a linear and norm-preserving isomorphism of $F_p^\delta \cap D$. In fact, h is the restriction of an orthogonal transformation in F_p^δ . Thus, h is a norm preserving map from $\partial D \cap F_p^\delta$ to $\partial D \cap F_p^\delta$ and the only possibility for h is either $h(a_j) = a_j$ or $h(a_j) = \sigma(a_j) = -a_j$, $j = 1, \dots, n$.

Recall that $a_j = \delta e_j$, the linearity of h implies either $h(e_j) = e_j$ or $h(e_j) = -e_j, \forall j$. Thus, h is either the identity map or the negative identity, in other words, $h = id$ or $h = \sigma$. The second case is not possible since then h is antiholomorphic rather than holomorphic. We conclude $h = id, K = id$ and $\text{Aut}(D) = L(G) \simeq G$. $\square$

REFERENCES

- [B-D] E. Bedford and J. Dadok, *Bounded domains with prescribed group of automorphisms*, Comment. Math. Helv. **62** (1987), 561-572.
- [C-G] C. Cheeger and D. Gromoll, *On the structure of complete manifolds of nonnegative curvature*, Ann. of Math. **96** (1972), 413-443.
- [C-M] S.-S. Chern and J. Moser, *Real hypersurfaces in complex manifolds*, Acta Math. **133** (1974), 219-271.
- [D-G] F. Docquier and H. Grauert, *Levisches problem und Rungescher Satz für Teilgebiete Steinscher mannigfaltigkeiten*, Math. Ann. **140** (1960), 94-123.
- [H-I] S. Halverscheid and A. Iannuzzi, *Maximal complexifications of certain homogeneous Riemannian manifolds*, Trans. Amer. Math. Soc. **355**, no. **11** (2003), 4581-4594.
- [HHK] P. Heinzner, A. Huckleberry and F. Kutzschebauch, *A real analytic version of Abels' theorem and complexifications of proper Lie group actions*, Lect. Notes Pure Appl. Math. **173** (1996), 229-273.
- [K] S.-J. Kan, *On rigidity of Grauert tubes over homogeneous Riemannian manifolds*, J. reine angew. Math. **577** (2004), 213-233.
- [S] N. Sibony, *A class of hyperbolic manifolds*, Recent developments in several complex variables, Ann. Math. Stud. **100** (1981), 357-372.
- [S-Z] R. Saerens and W. Zame, *The isometry groups of manifold and the automorphism groups of domains*, Trans. Amer. Math. Soc. **301**, no. **1** (1987), 413-429.
- [W] J. Winkelmann, *Realizing connected Lie groups as automorphism groups of complex manifolds*, Comment. Math. Helv. **79**, no. **2** (2004), 285-299.